


Fig. 1. Determination of methylation status of the MLH1 gene



Tab. 1. The mean age of patients from all risk groups, years

Age, years	Promoter hypermethylation of MLH1	
	Abs.	%
30-39	-	-
40-49	2	20.0
50-59	3	50.0
60-69	5	50.0
70-79	0	0

Tab. 2. The mean age of patients with Promoter hypermethylation of MLH1, years

Abstract #103 Figure 1/Table 1, 2

Disclosures None

#104 RISK FACTORS FOR SENTINEL LYMPH NODE METASTASIS IN ENDOMETRIAL CANCER (TRSGO-SLN-010)

¹Ibrahim Yalcin*, ²Salih Taskin, ³Ozguç Takmaz, ⁴Fuat Demirkiran, ³Mete Gungor, ⁵Nedim Tokgozoglu, ³Emine Karabuk, ⁴Tugan Bese, ²Duygu Altin, ⁶Hasan Turan, ⁷Ilker Kahramanoglu, ⁸Dogan Vatansever, ⁹Cetin Celik, ³Faruk Kose, ¹⁰Hamdullah Sozen, ¹⁰Samet Topuz, ⁴Macit Arvas, ²Firat Ortac, ¹¹Ali Ayhan, ⁸Cagatay Taskiran. ¹Ondokuz Mayıs University School of Medicine, Samsun, Turkey; ²Ankara University School of Medicine, Ankara, Turkey; ³Acibadem University School of Medicine, Istanbul, Turkey; ⁴Istanbul University Cerrahpasa School of Medicine, Istanbul, Turkey; ⁵Klinikum Nordfriesland, Husum, Germany; ⁶Mersin City Hospital, Mersin, Turkey; ⁷Private Clinic, Istanbul, Turkey; ⁸Koc University School of Medicine, Istanbul, Turkey; ⁹Selçuk University School of Medicine, Konya, Turkey; ¹⁰Istanbul University School of Medicine, Istanbul, Turkey; ¹¹Baskent University School of Medicine, Ankara, Turkey

10.1136/ijgc-2023-ESGO.273

Introduction/Background There is limited consensus on the optimal management approach for patients experiencing

mapping failure in endometrial cancer (EC). Understanding the risk factors that contribute to sentinel lymph node (SLN) metastasis is of paramount importance. This manuscript aims to provide a comprehensive analysis of the risk factors associated with SLN metastasis.

Methodology A total of 874 women with EC were included in this retrospective study. Out of the initial cohort of 874 patients, a total of 793 patients with successful SLN mapping were included and analysed to investigate the risk factors for SLN metastasis in EC.

Results SLN metastasis was detected in 73 (9.2%) patients. Among the metastatic cases, 20 (27.4%) patients had isolated tumour cells (ITC), 17 (23.3%) patients had micrometastasis, and 36 (49.3%) patients had macrometastasis in the sentinel lymph nodes. The results of the univariate analysis demonstrated a significant association between SLN metastasis and several factors, including age over 60 years, histology other than endometrioid, tumor grade 3, deep myometrial invasion, lymphovascular space invasion (LVSI), primary tumour diameter of 2 cm or larger, and cervical stromal invasion ($p < 0.05$). At the end of multivariate analysis, deep myometrial invasion [odds ratio (OR), 2.42; 95% confidence interval (CI), 1.29–4.56; $p = 0.006$], LVSI (OR, 7.27; 95% CI, 3.82–13.81; $p < 0.001$) and cervical stromal invasion (OR, 2.18; 95% CI, 1.13–4.21; $p = 0.020$) remained as independent risk factors for SLN involvement in women with EC.

Conclusion LVSI, deep myometrial invasion, and cervical stromal invasion emerged as independent risk factors for SLN metastasis in patients diagnosed with EC. In cases where the identified risk factors are absent, the omission of lymphadenectomy may be considered in instances of SLN mapping failure.

Disclosures The authors has no competing financial interests or conflicts of interest to disclose.

#110 ROLE OF EXTRACELLULAR VESICLES IN EARLY DIAGNOSIS OF ENDOMETRIAL CANCER

Irene Iavarone*, Antonella Reino, Maria Giovanna Vastarella, Rossella Moliterno, Maria Cristina Solazzo, Pasquale De Franciscis, Carlo Ronsini. *Department of Woman, Child and General and Specialized Surgery, University of Campania Luigi Vanvitelli, Naples, Italy*

10.1136/ijgc-2023-ESGO.274

Introduction/Background Extracellular vesicles are a class of cell-derived submicron particles, mediating cellular crosstalk through micro-RNA (miRNA). MiRNA are a group of RNA molecules, composed of 15–22 nucleotides each, post-transcriptionally regulating genes. Complementary mRNAs – into which miRNAs hybridise – are involved in implantation, tumour suppression, proliferation, angiogenesis, and metastatization defining tumour microenvironment. Despite endometrial biopsy being a standardized option to diagnose cellular atypia, non-invasive biomarkers may avoid discomfort of invasive procedures. The present study aims to evaluate distribution and regulation of differently expressed miRNAs (DEMs) in the context of endometrial cancer.

Methodology Following the recommendations in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement, we systematically searched PubMed, EMBASE, Scopus, Cochrane Library, and Science Direct databases in April 2023, adopting the string ‘Endometrial Neoplasms AND Exosomes’. We selected studies including patients with endometrial cancer, describing miRNA regulation in that