


Association between Patellar Instability and Increased Supratrochanteric Torsional Malalignment

Aytek Hüseyin Çeliksöz,^{*} MD, Yakup Alpay,[†] MD, Atakan Ezici,[†] MD, Tahir Koray Yozgatlı,[‡] MD, Ali Engin Daştan,[§] MD, Arman Vahabi,^{||} MD, and Baris Kocaoglu,^{‡¶} MD 

Investigation performed at Eskisehir City Hospital, Department of Orthopedics and Traumatology, Eskisehir, Turkey

Background: Recurrent patellar dislocation (RPD) is a common condition, particularly in adolescents, with several anatomic factors, such as increased femoral torsion, contributing to its development. Accurate assessment of these factors is essential for effective diagnosis and treatment planning.

Purpose: To examine the correlation between femoral torsion and RPD and to measure the level of torsion using magnetic resonance imaging (MRI).

Study Design: Cross-sectional study; Level of evidence, 3.

Methods: A total of 212 individuals were initially screened. Of these, 22 patients (31 femurs) who satisfied the inclusion criteria were included in the final analysis as the study group. The control group consisted of 28 patients (28 femurs) without a history of RPD. MRI measurements were conducted across 3 distinct axial sections. Supratrochanteric torsion (STT) value, infratrochanteric torsion (ITT) value, and femoral anteversion were analyzed utilizing T2-weighted turbo spin-echo sections. The distribution of the variables was determined using the Kolmogorov-Smirnov test. The independent-samples *t* test and chi-square test were used to analyze quantitative independent data. The Fisher exact test was used when the conditions appropriate for the chi-square test were not met.

Results: The mean femoral anteversion was higher in the patellar dislocation group, with a statistically significant difference (20.1 ± 3.8 vs 11.5 ± 1.8 ; $P = .0001$). The mean STT was significantly greater in the patellar dislocation group (41.3 ± 7.6 vs 35.2 ± 3.1 ; $P = .0002$). The mean ITT values were comparable between the groups (22.7 ± 2.8 vs 23.7 ± 2.3 ; $P = .14$).

Conclusion: Our results indicate that increased femoral anteversion in RPD originates from the supratrochanteric region of the femur. Future large-scale studies, specifically focusing on the relationship between femoral anatomic variables and patellar instability, could yield significant insights into this matter.

Keywords: recurrent patellar dislocation; femoral torsion; supratrochanteric torsion; MRI-based evaluation

Recurrent patellar dislocation (RPD) is a condition associated with the broader concept of patellar instability, which is primarily observed in adolescent patients.^{6,8} Female adolescents aged 10 to 17 appear to be the group with the highest risk.^{3,11} Despite the identification of factors such

as alterations in patellar height, trochlear dysplasia, and the rotational malalignment of both the femur (increased anteversion) and the tibia (lateralization of tibial tubercle) as important factors in the causation of RPD, the comprehensive etiology of this condition remains insufficiently delineated.^{7,14,17}

Recently, the influence of femoral axial malalignment on the etiology of various hip and knee pathologies garnered increased interest. Studies have established a correlation between increased femoral torsion and a range of conditions, including femoroacetabular impingement, anterior cruciate ligament injuries, and patellofemoral pain.¹⁹ Further studies have explored whether the specific

Correction (July 2025): This article has been updated to correct the affiliation for Baris Kocaoglu.

The Orthopaedic Journal of Sports Medicine, 13(7), 23259671251353754
DOI: 10.1177/23259671251353754
© The Author(s) 2025

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

location of the femoral torsion, supratrochanteric or diaphyseal, affects its association with these clinical presentations.^{1,10,12,13,17,18} Currently, there is no consensus on which location of the femoral torsion has a greater effect on the etiology of patellar instability.

In this study, we aimed to investigate the effect of supratrochanteric and diaphyseal femoral torsion on the etiology of RPD. Our hypothesis posits that femoral axial malalignment originating from the supratrochanteric region poses a higher risk for patellar instability than malalignment originating from the infratrochanteric region.

METHODS

Patient Selection

This retrospective age-matched case-control study was performed at the sports surgery department of an orthopaedic surgery clinic within a tertiary care center, between January 2021 and December 2023. Ethical approval for the study was granted by the local ethics committee. All cases diagnosed with RPD during the specified period were reviewed.

The inclusion criteria included patients with RPD, described as ≥ 1 patellar dislocation event, without multi-energy trauma, and patients aged < 18 years with complete medical and radiologic records. The exclusion criteria encompassed a history of fracture in the affected extremity, leg-length discrepancy, incomplete medical or radiologic records, and previous diagnosis of slipped capital femoral epiphysis, Legg-Calvé-Perthes disease, or connective tissue disorders such as Marfan syndrome or Ehlers-Danlos syndrome, as well as generalized joint hypermobility, which was defined as a Beighton score > 4 . Patients in the control group were selected by screening individuals who underwent femoral magnetic resonance imaging (MRI) at the same center during the same period, with matching based on age. In patients aged < 18 years, femoral neck angle variation between age groups has been well-documented; thus, age matching was performed by a range of ± 1 year.⁵ The control group, aged 12 to 18 years, had previously undergone bilateral femoral MRI scans for various reasons. Only patients without any history of patellar instability or related symptoms were included in the control group.

Starting in 2021, our center adopted a standardized protocol involving a whole-femur MRI to assess femoral

torsion in patients presenting with patellar instability. This transition to MRI for evaluating torsional malalignment was motivated by its efficacy in providing detailed anatomic assessments while avoiding radiation exposure.¹⁶

The study group: a total of 212 patients were initially screened for the study group, and 31 femurs of 22 patients who met the inclusion criteria were included in the final analysis. The control group included 28 patients (28 femurs) who had no history of patellar dislocation (Figure 1).

MRI Measurements

MRI measurements were conducted by 2 senior orthopaedic surgeons with > 5 years in the specialty (A.H.Ç., Y.A.), 2 weeks apart, each blinded to the previous results. After confirmation of sufficient intraclass correlation (> 0.8 ; good reliability), the mean values of each measurement were incorporated into the analyses.

MRI measurements were conducted across 3 distinct axial sections. For evaluations, T2-weighted turbo spin-echo sections were used. The first section was optimized for the sphericity of the femoral head, which involved drawing a line between the center of the circle placed on the femoral head to the center of the circle placed in the cervicobasilar junction at the femoral neck (first line). The second section was identified at the level where the lesser trochanter appeared most medially prominent, with a line drawn from the center of the femoral diaphysis to the apex of the lesser trochanter (second line). The third section was optimized to define the posterior condylar axis, which marks the most posterior aspect of the distal femoral articular surface (third line). The angular difference between the first and second lines was analyzed as the supratrochanteric torsion (STT) value, while the difference between the second and third lines was recorded as the infratrochanteric torsion (ITT) value (Figure 2). Femoral anteversion was analyzed as the angle between the first and third lines and was evaluated according to the method proposed by Fritz et al.¹⁰

Statistical Analysis

The mean, standard deviation, median lowest, median highest, frequency, and ratio values were used for the descriptive statistics of the data. The distribution of the variables was determined using the Kolmogorov-Smirnov

*Address correspondence to Baris Kocaoglu, MD, Acıbadem Üniversitesi Mehmet Ali Aydınlar Kampüsü Acıbadem Altunizade Hastanesi, İstanbul 34752, Turkey (email: bariskocaoglu@gmail.com).

*Koç University Hospital, Department of Orthopaedics and Traumatology, İstanbul, Turkey.

[†]Medical Park Maltepe Hospital, Department of Orthopaedics and Traumatology, İstanbul, Turkey.

[‡]Acıbadem University Faculty of Medicine, Department of Orthopaedics and Traumatology, İstanbul, Turkey.

[§]Department of Hand Surgery, İzmir City Hospital, İzmir, Turkey.

^{||}Ege University School of Medicine, Department of Orthopaedics and Traumatology, İzmir, Turkey.

Final revision submitted April 8, 2025; accepted May 1, 2025.

The authors declared that there are no conflicts of interest in the authorship and publication of this contribution. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval for this study was obtained from the local ethics committee (approval No. ESH/BAEK 2024/31).




Figure 1. Flowchart of the patient enrollment process.




Figure 2. (A) T1-weighted axial image identifying the center of the femoral head. (B) T1-weighted axial image identifying the center of the femoral head and femoral neck axis. Line (1) leads from the center of the circle placed on the femoral head to the center of the circle at the cervicobasilar junction of the femoral neck. (C) T1-weighted axial image identifying the section where the lesser trochanter appears most medially prominent. Line (2) indicates the detection of the lesser trochanteric apex with the largest medial extension. The angle between lines (1) and (2) describes supratrochanteric torsion. (D) T1-weighted axial image identifying the posterior condylar axis, which marks the most posterior aspect of the distal femoral articular surface. Line (3) shows the femoral posterior condylar axis. The angle between lines (2) and (3) describes infratrochanteric torsion.




Figure 3. The patellar dislocation group had high femoral anteversion (FA) and supratrochanteric torsion (STT), but no infratrochanteric torsion (ITT). There was a high association between femoral anteversion and high supratrochanteric torsion within the patellar dislocation group.

TABLE 1
Demographic Characteristics^a

Group	Patellar Dislocation Group (n = 31)	Control Group (n = 28)	P
Age, y	14.7 ± 1.8 (12-17)	14.5 ± 1.9 (12-17)	.68
Sex, female	17 (55)	16 (57)	.86
Side, right	16 (52)	14 (50)	.90

^aData are presented as mean ± SD or n (%).

TABLE 2
Comparison of Femoral Anteversion, STT, and ITT Between Groups^a

Group	Patellar Dislocation Group	Control Group	P
FA	20.1 ± 3.8 (15.6-24.8)	11.5 ± 1.8 (8.4-14.8)	< .001
STT	41.3 ± 7.6 (35.5-47.4)	35.2 ± 3.1 (30.9-39.2)	< .001
ITT	22.7 ± 2.8 (19.2-27.0)	23.7 ± 2.3 (19.1-27.5)	.14

^aData are presented as mean ± SD (range).FA, femoral anteversion; ITT, infratrochanteric torsion; STT, supratrochanteric torsion. P values compare differences between groups; P < .05 indicates statistical significance.

test. One-year age difference between groups was considered acceptable, and statistical comparisons were performed accordingly. The independent-samples *t* test and chi-square test were used to analyze quantitative independent data. The Fisher exact test was used when the conditions appropriate for the chi-square test were not met. The

SPSS Version 22.0 statistical program (IBM Corp) was used to perform all analyses. With a power of 80%, α level of .05, and 95% CI, the number of patients needed was calculated as 16 per group.

RESULTS

This study investigated data from 22 patients (31 femurs) with dislocation history and 28 control group patients with no dislocation history. The demographic characteristics were similar between the groups (Table 1). The mean number of dislocations was 2.5 in the patellar dislocation group (range, 1-5).

The mean femoral anteversion was greater in the patellar dislocation group (20.1 ± 3.8 vs 11.5 ± 1.8; $P = .0001$) (Table 2). The mean STT was significantly greater in the patellar dislocation group (41.3 ± 7.6 vs 35.2 ± 3.1; $P = .0002$). The mean ITT values were similar between the groups (22.7 ± 2.8 vs 23.7 ± 2.3; $P = .14$) (Figure 3).

In the patellar dislocation group, there was significantly higher femoral anteversion and STT. In contrast, the patellar dislocation group exhibited no significant ITT compared with controls (Table 2 and Figure 3).

DISCUSSION

The most significant finding was that increased femoral torsion originated from the supratrochanteric (41.3 ± 7.6 vs 35.2 ± 3.1; $P < .05$) region of the femur, rather than from the infratrochanteric (22.7 ± 2.8 vs 23.7 ± 2.3; $P = .14$) region or the distal femur when compared with normal femoral anteversion.

The study by Seitlinger et al,¹⁸ which initiated discussions regarding the anatomic site of torsional malalignment, had a notable effect; however, the broad age range of participants, spanning from 11 to 65 years old, is a limitation to consider. The study showed that femoral anteversion increases by approximately 1.5° per year, particularly in adolescents.⁵ Additionally, Archibald et al² stated that the torsional increase of the femur is slightly more related to the femoral neck than to the femoral shaft. In our study, we felt that femoral anteversion decreases with age by matching the control and study groups within the typical age range for femoral torsional abnormalities. We believe this approach enhanced the statistical reliability of the femoral anteversion angle measurements, which are known to vary with age.

Determining the anatomic site of the rotational deformity in the femur is critical to the treatment plan and clinical outcomes.^{4,9} Jia et al¹⁵ reported better results in patients having femoral torsion above the lesser trochanter in the treatment of RPD, although corrective osteotomies were performed distal to the femur regardless of the anatomic site of torsional malalignment. Our results suggested that the torsional malalignment in patients with RPD originated from supratrochanteric region rather than the infratrochanteric region. This raises the question of whether performing the osteotomy at this level eliminates the need for different soft tissue interventions or if the osteotomy level should be more proximal rather than distal. These results may inform future research comparing distal and proximal femoral derotation osteotomies in patients with RPD.

Although the importance of femoral torsion in RPD has been previously established, the exact location within the femur remained unclear, with the exception of the findings presented by Seitlinger et al.¹⁸ However, Fritz et al¹⁰ demonstrated that ITT or STT in femoroacetabular impingement may influence the treatment. The current study agrees with this for RPD.

Xu et al²⁰ studied segmental femoral torsion in patients with the RPD and concluded that midfemoral torsional malalignment was more correlated with RPD. While one of the limitations in the study by Xu et al was to pool patients from a broader range of age groups, their study reported important data on this debatable subject. Although the results of the current study appear to contradict those of Xu et al, we believe 2 key issues merit further investigation. First, torsional malalignment characteristics may vary across cohorts of different ethnic origins. Second, potential differences between the growing skeleton and the fully matured skeleton could influence these findings. Future studies focusing on more specific subgroups and homogeneous patient populations that can provide clearer answers to these questions would be valuable in this regard.

Limitations


This study has several limitations. First, it was retrospective and included a relatively small patient cohort, which

could have potentially biased the results. Second, the absence of data on other anatomic variables such as trochlear and tibial morphology, as well as the lack of matching for height and skeletal maturity, may have introduced selection bias. Although additional patellofemoral measurements such as tibial tuberosity–trochlear groove distance, Quadriceps angle (Q-angle), and sulcus angle were reviewed during patient screening, they were not included in the analysis, as the study focused specifically on the segmental origin of femoral torsion.

CONCLUSION

Our results indicate that increased femoral anteversion in RPD originates from the supratrochanteric region of the femur. Future large-scale studies, specifically focusing on the relationship between femoral anatomic variables and patellar instability, could yield significant insights into this matter.

ORCID iD

Baris Kocaoglu  <https://orcid.org/0000-0002-2537-0660>

REFERENCES

- Alpay Y, Ezici A, Kurk MB, Ozyalvac ON, Akpınar E, Bayhan AI. Increased femoral anteversion related to infratrochanteric femoral torsion is associated with ACL rupture. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(8):2567–2571. doi:10.1007/s00167-020-05874-0
- Archibald HD, Petro KF, Liu RW. An anatomic study on whether femoral version originates in the neck or the shaft. *J Pediatr Orthop.* 2019;39(1):e50–e53. doi:10.1097/BPO.0000000000001070
- Atkin DM, Fithian DC, Marangi KS, Stone ML, Dobson BE, Mendelsohn C. Characteristics of patients with primary acute lateral patellar dislocation and their recovery within the first 6 months of injury. *Am J Sports Med.* 2000;28(4):472–479. doi:10.1177/0363546500028040601
- Biedert RM. Reflections on rotational osteotomies around the patellofemoral joint. *J Clin Med.* 2021;10(3):474. doi:10.3390/jcm10030474
- Cibulka MT. Determination and significance of femoral neck anteversion. *Phys Ther.* 2004;84(6):550–558. doi:10.1093/ptj/84.6.550
- Danielsen O, Poulsen TA, Eystuoy NH, Mortensen ES, Hölmich P, Barfod KW. Familial association and epidemiological factors as risk factors for developing first time and recurrent patella dislocation: a systematic review and best knowledge synthesis of present literature. *Knee Surg Sports Traumatol Arthrosc.* 2023;31(9):3701–3733. doi:10.1007/s00167-022-07265-z
- Dejour H, Walch G, Nove-Josserand L, Guier Ch. Factors of patellar instability: an anatomic radiographic study. *Knee Surg Sports Traumatol Arthrosc.* 1994;2(1):19–26. doi:10.1007/BF01552649
- Dippmann C, Lavard P, Kourakis AH, et al. Good 5-year results and a low redislocation rate using an à la carte treatment algorithm for patellofemoral instability in patients with severe trochlea dysplasia. *Knee Surg Sports Traumatol Arthrosc.* 2025;33(2):401–412. doi:10.1002/ksa.12432
- Ferlic PW, Runer A, Seeber C, Thöni M, Seitlinger G, Liebensteiner MC. Segmental torsion assessment is a reliable method for in-depth analysis of femoral alignment in Computer Tomography. *Int Orthop.* 2018;42(6):1227–1231. doi:10.1007/s00264-017-3598-8
- Fritz B, Bensler S, Leunig M, Zingg PO, Pfirrmann CWA, Sutter R. MRI assessment of supra- and infratrochanteric femoral torsion:

- association with femoroacetabular impingement and hip dysplasia. *Am J Roentgenol*. 2018;211(1):155-161. doi:10.2214/AJR.17.18882
11. Gravesen KS, Kallemose T, Blønd L, Troelsen A, Barfod KW. High incidence of acute and recurrent patellar dislocations: a retrospective nationwide epidemiological study involving 24.154 primary dislocations. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(4):1204-1209. doi:10.1007/s00167-017-4594-7
 12. Hao K, Niu Y, Feng A, Wang F. Outcomes after derotational distal femoral osteotomy for recurrent patellar dislocations with increased femoral anteversion: a systematic review and meta-analysis. *Orthop J Sports Med*. 2023;11(7): 23259671231181601. doi:10.1177/23259671231181601
 13. Hao K, Niu Y, Kong L, Wang F. Medial patellofemoral ligament reconstruction combined with derotational distal femoral osteotomy yields better outcomes than isolated procedures in patellar dislocation with increased femoral anteversion. *Knee Surg Sports Traumatol Arthrosc*. 2023;31(7):2888-2896. doi:10.1007/s00167-022-07264-0
 14. İğrek S, Şahbat Y, Koç E, Çeliksöz AH, Topkar OM. Patellar lateralization, absence of joint hypermobility, and the mechanism of injury are associated with osteochondral fracture after first-time acute lateral patellar dislocation in adolescents: an MRI-based evaluation. *J Pediatr Orthop*. 2024;44(4):273-280. doi:10.1097/BPO.0000000000002629
 15. Jia Y, Bao H, Hou J, et al. Derotational distal femoral osteotomy yields better outcomes in patellar subluxation with proximal femoral torsion compared with distal femoral torsion: a retrospective comparative study. *J Orthop Surg Res*. 2024;19(1):640. doi:10.1186/s13018-024-05123-x
 16. Muhamad AR, Freitas JM, Bomar JD, Dwek J, Hosalkar HS. CT and MRI lower extremity torsional profile studies: measurement reproducibility. *J Child Orthop*. 2012;6(5):391-396. doi:10.1007/s11832-012-0434-y
 17. Perry AK, Maheshwer B, DeFroda SF, et al. Patellar instability. *JBJS Rev*. 2022;10(11):e22.00082. doi:10.2106/JBJS.RVW.22.00054
 18. Seitlinger G, Moroder P, Scheurecker G, Hofmann S, Grelsamer RP. The contribution of different femur segments to overall femoral torsion. *Am J Sports Med*. 2016;44(7):1796-1800. doi:10.1177/0363546516639945
 19. Teitge RA. Pathological torsion of the lower limb. In: LaPrade RF, Chahla J, eds. *Evidence-Based Management of Complex Knee Injuries: Restoring the Anatomy to Achieve Best Outcomes*. Elsevier; 2020:336-342. doi:10.1016/B978-0-323-71310-8.00029-3
 20. Xu Z, Song Y, Deng R, Zhang Z, Wang H, Yu JK. Pathological thresholds of segmental femoral torsion in patients with patellar dislocation: influence on patellofemoral malalignment. *Orthop J Sports Med*. 2022;10(10):23259671221125218. doi:10.1177/23259671221125218